



Full Length Research Paper

Pesticidal plants used by farmers converting to organic farming in Uganda

Aggrey Oryema Nuwagaba

Faculty of Agriculture, Uganda Martyrs University, Nkozi, P. O. Box 5498 Kampala, Uganda.

Accepted 29 July, 2016

Use of synthetic pesticides in developing countries is not only limited by their being expensive but also the small (uneconomic) fields whose limited production costs cannot offset costs of agricultural implements like agro-chemicals. Subsistence farmers, therefore, have no choice but to use local methods of controlling pests, one of which is the use of traditional and of late introduced pesticidal plants' extracts. In this study, whose main objective was to record all pesticidal plants used in Southern Uganda, Masaka district, it was established that thirty four species belonging to eighteen families are currently used in traditional plant production. Most useful species were *Azadirachta indica* and *Tagetes minuta* while the most frequently cited families were Meliaceae and Euphorbiaceae. It was noted that of the plant species recorded, some plants like *A. indica*, *Melia azedarach*, and *T. minuta* are already scientifically established pesticidal plants whereas others like *Euphorbia tirucalli*, *Bidens pilosa*, *Vernonia amygdalina* may be known for other uses but not for this purpose and hence the need for their efficacy evaluation. Some important pesticidal plants like *Abrus precatorius*, *Euphorbia candelabrum* and *Phoenix reclinata* were reportedly becoming increasingly rare and would need conservation. The need to carry out such surveys in order to obtain inventories was observed and recording this knowledge before it disappears with the aging farmers was seen as urgent.

Key words: Indigenous knowledge, inventory, developing countries, conservation.

INTRODUCTION

In most developing countries, use of modern synthetic pesticides is limited and sometimes non-existent (Scialabba, 2000). This is so not only because they are expensive but also because of the small fields cultivated by subsistence farmers making the use of such pesticides uneconomic. Yet, most of these countries lie in tropical and subtropical areas where pests and diseases are abundant throughout the year.

The implication to this is that in these areas, pests and diseases pose a major problem in agricultural production. According to Oerke and Dehne (2004), pests contribute 30-40% of crop loss worldwide while the loss in the tropics is reported to be even higher than 40% (FAO,

2003). These figures show that the battle against pests is a difficult one even with the advanced technology that exists today.

In developing countries, where technological advancement is still low and the use of modern methods is subjacent, the most credible alternative appears to be the use of traditional/cultural methodologies (Pei, 2001; Rates, 2001; Muhammad and Awaisu, 2008) - one of which is the use of traditionally-known pesticidal plants as a remedy for pest infestations.

The use of botanicals for pest control is as old as agriculture itself. Thacker (2002) reports the use of

*Corresponding author: E-mail: oryema2016@gmail.com

Author(s) agreed that this article remain permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

tobacco (*Nicotiana tabacum* L.) leaves for fumigation in stores as early as the late 1500s, the use of *Sabadilla officinale* A. Grey ex Benth. for crop protection in the mid 1500s, and application of *Quassia* spp extracts against aphids during the early 1600s among others.

While these botanicals were used as early as reported above, the discovery of synthetic pesticides in the early 1900s tended to overwhelm their use because of the many advantages synthetic pesticides appeared to have at that time. For example, DDT was reported to have a knockdown effect on most insects, high persistence in the environment, ease of application together with having a broad spectrum (DeLong, 1948; Walker, 2000) which advantages the botanicals appeared not to possess.

It was only after environmentalists like Carson (1962) started realizing that these advantages were in effect potential disadvantages that scientists started to question the future prospects of synthetic pesticides. Therefore, botanicals are not new on the pest management scene but have recently been re-discovered as a route through which to escape the many disadvantages of synthetic pesticides.

However, as Asman (2008) and Muhammad and Awaisu (2008) say, only a few are presently in commercial use. Therefore, there is need to explore, document and evaluate the use of more plants. Yet, Ankli et al. (1999) and Gradé (2008) note that knowledge of such herbs is usually in the hands of a small group of people who guard it jealously for their advantage. In many cases, it is passed on between generations by word of mouth or observing elders' activities and is often not documented. In many developing countries like Uganda, documentation and scientific evaluation of medicinal plants in general and botanicals for pest management in particular is still wanting. Available literature reveals that some ethnobotanical survey have been made in Uganda for traditional plants used in human medicine (Freiburghaus et al., 1996; Kamatenesi-Mugisha et al., 2007a; Katuura et al., 2007; Ssegawa and Kasenene, 2007; Waako et al., 2007), veterinary medicine (Tabuti et al., 2003; Bukonya-Ziraba and Kamoga, 2007; Gradé et al., 2007; Katuura et al., 2007), cultural values (Kakudidi, 2004a; Kakudidi, 2004b), but only limited amount of information concerning pesticidal/insecticidal plants used in this region is available (Mugisha-Kamatenesi et al., 2008). This is in stark contrast with the ideal requirements of the ongoing campaigns to 'go organic' that are slowly but steadily catching up in this region. Organic farming by principle discourages the use of synthetic agrochemicals and promotes the use of ecological techniques like use of plant-based pest remedies.

Therefore, this study was set up to contribute to documentation of pesticidal plants used in Uganda especially in the agricultural district of Masaka (south Uganda) where many farmers are converting to organic farming. The study objectives were: a) documentation of all known pesticidal plants in the area; b) establishment

of pests on which these pesticidal plants are applied; and c) finding out their mode of application/formulation. Results will help to obtain a record of these plants from the farmers especially those who are elderly before they die/disappear with their knowledge and will stimulate further research on several aspects like conservation of the species and efficacy evaluation of plant substances against the pests they are claimed to control.

MATERIALS AND METHODS

Study area

The study was carried out in Masaka district located between 31° 12' and 32° 06'E and 0° 48' and 1° 20' S in South Uganda (Figure 1). Found on the shores of Lake Victoria, Masaka district is one of the most important agricultural areas in Uganda due to its favorable climatic conditions. The district has a bimodal type of rainfall with an annual average of 1200 mm and mild equatorial temperatures ranging between 22 and 26°C. Following the bimodal type of rainfall, the district has two growing seasons, March to June and October to December that support the growing of crops the whole year around. Main crops grown include bananas (*Musa* spp), beans (*Phaseolus vulgaris* L.), cassava (*Manihot esculenta* Crantz.), maize (*Zea mays* L.), coffee (*Coffea* spp) and a range of other (tropical) vegetables, fruits and cereals (Tenywa et al., 1999).

Data collection

The study was carried out with full knowledge and support of local councils (LCs) of the area that constitute the local administration at different levels of the district. Preparatory stages of the study involved holding meetings with local leaders who eventually introduced the research team to farmers. On the advice of district agricultural staff, it was found appropriate to include one agricultural field staff on the team to help with introduction and identification of 'right' respondents. Therefore, the survey team included an agricultural extension worker and a field officer recruited from the area, who also doubled as a translator where it was deemed necessary. The survey was conducted in the local language (Luganda) except for educated respondents who chose to use English.

During data collection, we opted for open-ended interviews in order to obtain quantitative data with a qualitative depth, devoid of researcher input. Mossholder et al. (1995) say that open-ended interviews offer a way of providing qualitative depth in a survey-based research and have an advantage of allowing respondents answer in their own frames of reference, implicitly reducing the prime influence of leading questions following researcher suggestions. We also used direct observation to record observable data on plant characteristics as recommended by Etkins (1993).

The survey was carried out between October 2007 and March 2008 and five sub-counties including Bigasa (0° 04' 59" S; 31° 38' 00" E), Bukoto (0° 23' 09" S; 31° 37' 38" E), Kitanda (0° 44' 59" S; 31° 35' 58" E), Lwengo (0° 24' 58" S; 31° 24' 29" E) and Buwunga (0° 22' 22" S; 31° 47' 35" E) were assessed (GPS coordinates taken are for farms of lead farmers in the sub-county). These sub-counties were chosen on advice of Masaka agricultural staff and using the criterion that they were among the leading areas in organic farming adoption; which implies extensive use of botanicals for pest management.

In each sub-county, twenty-five farmers, (about 1/3 of all farmers in a sub-county) and one extension staff were interviewed making a




Figure 1. Map of Uganda districts showing the study area (Masaka district). Adapted from Google maps.

total number of 130 respondents in the whole district. All respondents were adults (70 males and 60 females) between ages of 19 and 70 of African descent that is, about 90% were baganda, 6% were banyankole, 5% were of Rwandese origin and 1% were of luo origin.

During the survey, each respondent was requested to give information pertaining to the plants' local name, plant part used, method of formulation/application mode of part used and pest(s) which the latter was used against. Finally, voucher specimens (JM 1-34) were prepared from each plant mentioned and deposited at the herbarium of Makerere University. Plants were identified by comparison with authentic herbarium specimens and sometimes with assistance from experts at Makerere University herbarium.

Data analysis

All data was recorded in previously designed data sheets to reflect

different objectives. For analysis, species and families recorded were assessed for User Value (UV) (Aburjai et al., 2007; Heinrich et al., 1998) - a quantitative method that demonstrates the relative

$$\text{importance of species locally } UV = \frac{\sum U}{n},$$

where UV is the user value, U is the number of user citations and n is the number of respondents.

Informant consensus ICF (Aburjai et al., 2007; Ankli et al., 1999) was calculated to indicate information homogeneity. According to the latter authors, ICF will be lower (closer to 0), if there is a large variation in plant use or when users do not exchange information about plant uses. High values (close to 1) reflect well-defined plant use or information exchange between respondents. All calculations and graphic presentations were carried out in Microsoft office Excel 2007. UV and ICF values are reflected in Tables 1 and 3.

Table 1. Record of species, their families, plant parts used and Use Values (UVs).

Species name and Voucher no.	Local name	Family	Part used	Times mentioned	UV
<i>Abrus precatorius</i> L. JM-001	Lusiiti	Fabaceae	L, S	23	0.176
<i>Allium sativum</i> L. JM-002	Katungulucumu	Alliaceae	L	56	0.430
<i>Annona senegalensis</i> L. JM-003	Kisitaferi	Annonaceae	L,B	55	0.423
<i>Artemisia annua</i> L. LM-004	Artemisia	Asteraceae	L,FI	67	0.515
<i>Asparagus africanus</i> Lam. JM-005	Kadaali	Asparagaceae	L(spines)	45	0.346
<i>Azadirachta indica</i> A.Juss JM-006	Neem	Meliaceae	L,B R,F	125	0.961
<i>Bidens pilosa</i> L. JM-007	Ssere	Asteraceae	L	36	0.276
<i>Cannabis sativa</i> L. JM-008	Njagga	Cannabaceae	L, S, F	52	0.400
<i>Capsicum frutescens</i> L. JM-009	Kamulari	Solanaceae	F	87	0.669
<i>Carica papaya</i> L. JM-010	Mupaapali	Caricaceae	R,B	32	0.246
<i>Chrysanthemum coccineum</i> Willd. JM-011	Pyrethrum	Asteraceae	L, FI	23	0.176
<i>Citrus aurantifolia</i> Swingle JM-012	Nimawa	Rutaceae	F,L	32	0.246
<i>Cupressus lusitanica</i> L. JM-013	Kapripusi	Cupressaceae	L,B	54	0.415
<i>Cupressus sempervirens</i> L. JM-014	Ssedero	Cupressaceae	S,L	60	0.461
<i>Cymbopogon nardus</i> L.(Rendle) JM-015	Mutete	Poaceae	L	45	0.346
<i>Datura stramonium</i> L. JM-016	Ruziringa	Solanaceae	L	68	0.523
<i>Eucalyptus globulus</i> Labill. JM-017	Kalitunsi	Myrtaceae	L,B	60	0.461
<i>Eucalyptus grandis</i> W.Hill.ex Maid JM-018	Kalitunsi	Myrtaceae	L,B	56	0.430
<i>Euphorbia candelabrum</i> Tremaut. ex Kotschy JM-019	Nkukuulu	Euphorbiaceae	Latex, B, R Latex, B,	49	0.376
<i>Euphorbia tirucalli</i> L. JM-020	Nkoni	Euphorbiaceae	ash	46	0.353
<i>Jatropha curcas</i> L. JM-021	Kiryowa	Euphorbiaceae	Sap, F,S, B	103	0.792
<i>Lantana camara</i> L. JM-022	Kayukiyuki	Verbenaceae	L	69	0.530
<i>Melia azedarach</i> L. JM-023	Lira	Meliaceae	L, R,B	110	0.846
<i>Mucuna pruriens</i> Bak. JM-024	Mucuna	Fabaceae	L,R	32	0.246
<i>Nicotiana tabacum</i> L. JM-025	Taaba	Solanaceae	L	98	0.753
<i>Phoenix reclinata</i> Jacq. JM-026	Mukindo	Arecaceae	Sap, ash	15	0.115
<i>Phytolacca dodecandra</i> L'Herit JM-027	Luwoko	Phytolacceae	L, F	43	0.331
Plant ash N/A	Evuu	N/A	Ash	101	0.777
<i>Ricinus communis</i> L. JM-028	Nsogasoga	Euphorbiaceae	S	87	0.669
<i>Schinus molle</i> L. JM-029	Kishenda	Anacardiaceae	L, F	65	0.500
<i>Solanum lycopersicum</i> L. JM-030	Enyaanya	Solanaceae	F	24	0.184
<i>Tagetes minuta</i> L. JM-031	Kawunyira	Asteraceae	L	118	0.907
<i>Tephrosia vogelii</i> Hook.f. JM-032	Muluku	Fabaceae	L	88	0.676
<i>Tithonia diversifolia</i> (Hehsl.) A. Gray JM-033	Ekimyula	Asteraceae	F,L	75	0.577
<i>Vernonia amygdalina</i> Del. JM-034	Omululuza	Asteraceae	L	35	0.269

Key: L , Leaf; B, Bark; F, Fruit; S, Seed; R, Root.

RESULTS

Record of species

Table 1 shows the inventory of all species recorded from the survey. Thirty-four species belonging to 18 families were recorded. The most important pesticidal species were *Azadirachta indica*, *Tagetes minuta*, *Melia* 0.85 and from 0.7 to 0.12 with *Phoenix reclinata* (0.12), *Abrus*

precatorius (0.18) and *Chrysanthemum coccineum* (0.18) scoring the least (Table 1).

The most used families were Meliaceae, Euphorbiaceae, Solanaceae and Verbenaceae with UVs of 0.90, 0.55, 0.53 and 0.53, respectively while Arecaceae 0.79, respectively. Other species scored UVs ranging *azedarach* and *Jatropha carcus* with UVs of 0.96, 0.91, (0.12) and Rutaceae (0.25) were indicated as the least useful (Table 3). Only two families namely Poaceae

Table 2. Record of pesticide mode of formulation and pests they control.

Species name	Mode of formulation	¹Pest/ disease/deficiency treated
<i>Abrus precatorius</i>	Water extract	Worms
<i>Allium sativum</i>	Trap crop, water extract	Field pests, storage pests
<i>Annona senegalensis</i>	Water extract	Fungicidal properties, insects
<i>Artemisia annua</i>	Water extract	Mosquitoes, flies
<i>Asparagus africanus</i>	Physical trap/thorns	Birds, moths, bats
<i>Azadirachta indica</i>	Water extract/crashed seeds	Most insects, mosquitoes
<i>Bidens pilosa</i>	Water extract	Aphids
<i>Cannabis sativa</i>	Water extract, smoke, trap crop	Storage pests, insect pests, coccidiosis, anti-biotic
<i>Capsicum frutescens</i>	Water extract, crashed seeds	Cut worms, ants, snails, storage pests
<i>Carica papaya</i>	Leaf water extract, crashed seeds	Blight, animal worms
<i>Chrysanthemum coccineum</i>	Oil extract , water extract	Most pests
<i>Citrus aurantifolia</i>	Water extract, trap crop	Insect pests
<i>Cupressus lusitanica, Sempervirens</i>	Physical admixture, water extract	Storage pests, houseflies
<i>Cymbopogon nardus</i>	Trap crop, oil extract	Lepidoptera pests, beetles, aphids
<i>Datura stramonium</i>	Water extract	Insects
<i>Eucalyptus globulus</i>	Water extract, physical admixture, oil extract	Storage pests, repellent
<i>Eucalyptus grandis</i>	Water extract, physical admixture, oil extract	Storage pests, repellent
<i>Euphorbia candelabrum</i>	Latex spray, water extract	Termites, cutworms
<i>Euphorbia tirucalli</i>	Latex spray, ash dusting, physical admixture	Aphids, safari ants, cutworms
<i>Jatropha curcas</i>	Latex spray, water extract(seeds)/ oil	Insect pests, animal worms
<i>Lantana camara</i>	Water extract, physical mixture	Insect pests, storage pests
<i>Melia azedarach</i>	Water extract	Most insects, worms
<i>Mucuna pruriens</i>	Water extract, intercrop	Nitrogen deficiency
<i>Nicotiana tabacum</i>	Water extract, smoke, physical admixture	Storage pests, soil pests ,domestic pests, repels snakes
<i>Phoenix reclinata</i>	Ash/dusting	Storage pests
<i>Phytolacca dodecandra</i>	Water extract	Snails, insect pests ,fungi
Plant ash	Dusting, water mixture	Cut worms, banana weevil, storage pests, fungi
<i>Ricinus communis</i>	Water extract, crushed seed /oil	Storage pests, domestic pests, animal worms
<i>Schinus molle</i>	Water extract, crushed seeds	Worms, storage pests
<i>Solanum lycopersicum</i>	Water extract, repellent crop	Aphids, thrips, weevils
<i>Tagetes minuta</i>	Water extract, repellent crop	Most insects, nematodes
<i>Tephrosia vogelii</i>	Water extract	Insect pests, ticks, fungi, mites, moles
<i>Tithonia rotundifolia</i>	Trap crop	Nematode trap plant
<i>Vernonia amygdalina</i>	Water extract	Insect pests, malaria parasites, worms

¹Pests/diseases controlled were grouped into four broad uses: field pests, storage pests, domestic pests, veterinary pests and others/unclassified for calculation of ICF whose values were above 0.9 in all categories.

and Arecaceae belong to monocotyledonous subdivision while the rest are dicotyledonous.

Plant parts used as source of pesticide

Leaves were named as plant parts most-used in formulation of pesticides. They were reported for 26 species out of the total 34 cited in the survey. They were

followed by seeds/fruits and the bark with 11 and 10 species, respectively.

Ash made by burning wood and flowers were the least with 2 species each (Figure 3). However, ash was generally mentioned as a constituent of many concoctions. Several intersections were observed where some plants were named for more than one part. *A. indica*, *M. azedarach*, *Cannabis sativa* and *J. curcas* were named for using almost all parts while others were




Figure 2. Mode of formulation of pesticides among species recorded.




Figure 3. Plant parts used as source of pesticide.

named for one or two parts (Table 1).

Formulation/mode of utilization of plant parts

Water extract was the most commonly named mode of formulation accounting for nearly half (49%) of all formulations. Others included use of the whole plant as an intercrop that is, trap crop/repellent (10%), physical admixture of plant part with produce (10%), use of crushed seed cake (8%), application of plant oil extract (8%), latex spray (5%), ash admixture (5%) use of thorns as deterrents (2%), and use of smoke from burning plant parts (3%) (Figure 2).

Most plants were reported to have several ways in which pesticides could be formulated from them. For example *J. curcas*, *E. tirucalli*, *Eucalyptus* spp, *Nicotiana tabacum* and *C. sativa* were named for three ways, while the majority was reported for two ways and a few others for only one (Table 2).

Pests the plants are used against

Numerous pests were mentioned during the survey but it was apparent that farmers were neither familiar with formal classification nor names of pests and diseases. Most farmers gave broad answers such as weevils, storage pests, caterpillars, insects, moths or field pests. It was therefore difficult to obtain meaningful data for comparison.

During interviews, respondents also pointed out, the difficulty of naming particular pests managed by certain plants because extracts are used when there is infestation in the field without establishing particular pests being controlled. After all, many plants are used in combination with others. Therefore, plants were reported to be used against a range of pests. For analysis, pest data collected were organized in broad groups namely: field pests, storage pests, veterinary pests, domestic pests and others which could not be classified. All groups include the corresponding diseases. Table 3 shows that the most-cited pests were field pests named for 29 plant species in 1,270 user reports, while veterinary pests scored least with 5 species in 326 reports. However, all groups returned a high ICF that is, above 0.9 indicative of the fact that there is a high user consensus among the farmers and a likelihood of sharing ideas about use of botanicals.

DISCUSSION

Ethnobotanical surveys are made for different reasons. Some of these include assessment of functions of plants, for example identification of medicinal species (Jouad et al., 2001; Kamatenesi-Mugisha et al., 2007b; Katuura et al., 2007) and analysis of species diversity in a given area (Oryemoriga et al., 1995). Other reasons include determination of species conservation status (Schemske et al., 1994; van Jaarsveld et al., 1998) or when trying to identify new plant species in an area that is not yet extensively studied (Oryemoriga et al., 1995).

This study was meant to establish a record of pesticidal plant species used in Masaka district, South Uganda and how they are utilized.

It was found out that 34 species distributed in 18 families are being used in Masaka district as pesticidal plants (Table 1). Some of these species are already established /confirmed pesticidal plants such as *A. indica*, *M. azedarach*, *J. curcas*, *Tagetes. minuta*, *Tithonia rotundifolia*, *Chrysanthemum* spp (Isman, 2006) but there

Table 3. Family use values.

Family	No. of species	% of all species	Use citations	% use citations	UV
Asteraceae	6	17	354	17.41	0.453
Solanaceae	4	12	277	13.62	0.532
Euphorbiaceae	4	12	285	14.01	0.548
Fabaceae	3	9	143	7.03	0.366
Cupressaceae	2	6	114	5.60	0.438
Meliaceae	2	6	235	11.55	0.903
Myrtaceae	2	6	116	5.70	0.446
Alliaceae	1	2.9	56	2.75	0.430
Anacardiaceae	1	2.9	65	3.19	0.500
Annonaceae	1	2.9	55	2.75	0.423
Arecaceae	1	2.9	15	0.73	0.115
Asparagaceae	1	2.9	45	2.21	0.346
Cannabaceae	1	2.9	52	2.55	0.400
Caricaceae	1	2.9	32	1.57	0.246
Phytolacceae	1	2.9	43	2.11	0.330
Poaceae	1	2.9	45	2.21	0.346
Rutaceae	1	2.9	32	1.57	0.246
Verbenaceae	1	2.9	69	3.39	0.530

are some species like *E. tirucalli*, *Euphorbia candelabrum*, *Bidens pilosa* and *Vernonia amygdalina* that would require efficacy evaluation and publication/popularization for use as pesticidal plants. Therefore, there is a need to substantiate findings of this study by carrying out efficacy studies and other related work that may lead to confirmation and recommendation of such plants for more extensive use.

Meliaceae and Euphorbiaceae were reported to be the most useful families in this area for having a good number of species with pesticidal features. This means that they are good families to start from when looking for species of pesticidal importance.

Plants produce a variety of secondary metabolites to protect themselves against pathogens and herbivores and/or to influence the growth of neighboring plants (Fraenkel, 1969; Swain, 1977; Edwards, 1992; Sirikantaramas et al., 2008). The same authors go on to say that some of these metabolites are toxic to the producing cells when their target sites are present in the producing organisms and must therefore be demobilized. Such substances include alkaloids (Dethier, 1980), tannins (Bernays, 1981), phenols (Palo, 1984), terpins (Schutte, 1984), and according to Gatehouse (2002), these are the substances man can exploit to make pesticides of botanical origin for pest control.

Our results are indicative of the fact that leaves constitute a large portion of the said secondary substances. This conclusion was reached at by considering our results (Figure 3) which show that most-exploited tissues of pesticidal plants are leaves (76%) followed by fruits and seeds (53%) and then the

bark with 29%. These findings could be attributed to the positioning of plant parts on a plant. Plant defense theories suggest that chemical or structural defences should be maximized when and where browsing is most likely to occur (Massei et al., 2000). Leaves are exposed and conspicuous, which makes them easy targets for herbivore attack. It is not surprising, therefore, that plants tend to deposit and localize secondary substances in exposed parts such as leaves and fruit/flowers to act as deterrents to herbivores. Plants without conspicuous leaves like *Euphorbia* spp. utilize their green stems/latex for such a purpose.

Similar findings have been reported by Kamatenesi-Mugisha et al. (2007a), Maregesi et al. (2007), Ssegawa and Kasenene (2007) giving a clue to researchers on the cardinal parts to assess when in search of pesticides from plants.

Crops in tropical countries are heavily attacked by a variety of pests and diseases which are active throughout the whole year, probably due to tropical conditions (Kamatenesi-Mugisha et al., 2007a). Farmers, therefore have to fight these pests in order to obtain tangible outputs from fields. However, during the present study, farmers could not point out exactly the particular pests managed by the mentioned pesticidal plants which could be a limitation for pest management. They reported pests generally as storage pests, insects, weevils, field pests and others (Table 2). Such type of identification is incoherent and groups pests in categories that cannot help to draw scientific conclusions. While this could be attributed to low literacy rates of the farmers in this region (Dent, 2007), it could also be due to the fact that a

remedial concoction is often composed of a number of plant species and targets several pest species in the field or in storage (Kamoga per.comm). The need for research to elucidate chemical composition of the medicinal species and evaluate specific pests managed by particular plants species through efficacy studies was observed.

During our analysis, it was found appropriate to categorize the mentioned pests in broad groups namely field pests, domestic pests, veterinary pests and storage pests in order to give a broad view of pests managed by these plants. According to this categorization, field pests were the most-cited with 1270 user reports on 29 species, followed by storage pests (661 user reports on 16 species) indicative of the dominant type of occupation in the area studied. Masaka district is dominated by arable farming with little emphasis on pastoralism (Tenywa et al., 1999). It is likely that different results would have been obtained in a predominantly pastoral region (Gradé, 2008).

During interviews with extension staff, it was revealed that certain medicinal plants are no longer available in the area studied and farmers have to travel long distances to harvest them. Species named include: *Abrus precatorius*, *Phoenix reclinata*, and *E. candulubrum*, which are mainly woodland species. Without substantiation, farmers also reported that some plants are completely unavailable and cannot be found anywhere, which could have implications of extinction. Most of South Uganda and Masaka district in particular was originally a woodland area where these trees were once abundant (Hamilton, 1974). However, most of them have long been cut down for timber and while opening up agricultural land as the population increased. This is not surprising because over the years, an increasing number of plants have been red-listed as threatened species by IUCN in Uganda. For example, in 1997, IUCN had a red-list of 15 plants in Uganda (Walter and Gillet, 1999), which increased to 33 in 2002 (Earthtrends, 2003) and to 38 in 2008 (IUCN, 2009). This confirms our and other people's fears e.g. Hedberg (1993), Cox (2000) that unless these plants and the traditional knowledge thereof are documented now, they will soon face extinction. Therefore, due considerations to conserve these species (for example by domestication) should be made as quickly as possible before they disappear completely.

Conclusion

This study has shown that numerous plant species are used in this region for purposes of pest management. Notable ones such as *A. indica*, *M. azedarach*, and *T. minuta* dominate the application scene but a few 'new ones' like *B. pilosa* and *E. tirucalli* were also documented (for this purpose) for the first time in this region.

Therefore, there is need to establish their efficacy and

identify the pests against which their extracts are most active. Also, the need for conservation of such species of pesticidal importance was noted. The earlier it is done, the better for pest management and biodiversity.

REFERENCES

- Aburjai T, Hudaib M, Tayyem R, Yousef M, Qishawi M (2007). Ethnopharmacological survey of medicinal herbs in Jordan, the Ajloun Heights region. *J. Ethnopharmacol.*, 110: 294-304.
- Ankli A, Sticher O, Heinrich M (1999). Medical ethnobotany of the Yucatec Maya: Healers' consensus as a quantitative criterion. *Econ. Bot.*, 53(2): 144-160.
- Bernays EA (1981). Plant tannins and insect herbivores: an appraisal. *Ecol. Entomol.*, 6(4): 353-360.
- Bukenya-Ziraba R, Kamoga D (2007). An inventory of medicinal plants used in treating poultry diseases in Jinja district, eastern Uganda. *Afr. J. Ecol.*, 45: 31-38.
- Carson R (1962). The silent spring. <https://www.rachealcarson.org>. Accessed on 23 April 2008.
- Cox PA (2000). Will tribal knowledge survive the millennium? *Sci.*, 287(5450): 44-45.
- DeLong M (1948). DDT and the Insect Problem. *Q. Rev. Biol.*, 23(2): 159-160.
- Dethier VG (1980). Evolution of receptor sensitivity to secondary plant substances with special reference to deterrents. *Am. Nat.*, 115(1): 45.
- Earthtrends (2003). Biodiversity and protected areas in Uganda. www.earthtrends.wri.org. Accessed on 20 June 2009.
- Edwards PJ (1992). Resistance and defense: The role of secondary plant substances. *Environ. Plant Boil. Ser.*, 23: 69-84.
- Etkins NL (1993). Anthropological methods in ethnopharmacology. *J. Ethnopharmacol.*, 38: 93-104.
- FAO (2003). Crop protection in the context of Agricultural development. Report No. 232/2003.
- Fraenkel G (1969). Evaluation of our thoughts on secondary plant substances. *Entomol. Exp. Appl.*, 12(5): 473-486.
- Freiburghaus F, Kaminsky R, Nkunya MHH, Brun R (1996). Evaluation of African medicinal plants for their in vitro trypanocidal activity. *J. Ethnopharmacol.*, 55(1): 1-11.
- Gatehouse JA (2002). Plant resistance towards insect herbivores: a dynamic interaction. *New Phytol.*, 156(2): 145-169.
- Gradé JT (2008). Ethnoveterinary knowledge in pastoral Karamoja, Northern Uganda. Ph.D. thesis. University of Ghent, p. 218.
- Gradé JT, Tabuti JRS, Van Damme P, Arble BL (2007). Deworming efficacy of *Albizia anthelmintica* in Uganda: preliminary findings. *Afr. J. Ecol.*, 45(3): 18-20.
- Dent VF (2007). Local economic development in Uganda and the connection to rural community libraries and literacy. *New Libri. World* 108(5/6): 203-217.
- Hamilton A (1974). Distribution patterns of forest trees in Uganda and their historical significance. *Plant Ecol.*, 29: 21-35.
- Hedberg I (1993). Botanical methods in ethnopharmacology and the need for conservation of medicinal plants. *J. Ethnopharmacol.*, 38(2-3): 121-128.
- Heinrich M, Ankli A, Frei B, Weimann C, Sticher O (1998). Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc. Sci. Med.*, 47: 1859-1871.
- Isman MB (2008). Botanical insecticides: for richer, for poorer. *Pest Manage. Sci.*, 64: 8-11.
- Isman MB (2006). Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu. Rev. Entomol.*, 51: 45-66.
- IUCN (2009). IUCN 2008 red list of threatened species. www.iucnredlist.org. Accessed on 15 July 2009.
- Jouad H, Haloui M, Rhiouani H, El Hilaly J, Eddouks M (2001). Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). *J. Ethnopharmacol.*, 77(2-3): 175-182.
- Kakudidi EK (2004). Cultural and social uses of plants from and around

- Kibale National Park, Western Uganda. *Afr. J. Ecol.*, 42: 114-118.
- Kakudidi EK (2004). Folk plant classification by communities around Kibale National Park, western Uganda. *Afr. J. Ecol.*, 42: 57-63.
- Kamatanesi-Mugisha M, Oryem-Origa H, Olwa O (2007). Medicinal plants used in some gynaecological morbidity ailments in western Uganda. *Afr. J. Ecol.*, 45: 34-40.
- Katuura E, Waako P, Tabuti JRS, Bukenya-Ziraba R, Ogwal-Okeng J (2007). Antiplasmodial activity of extracts of selected medicinal plants used by local communities in western Uganda for treatment of malaria. *Afr. J. Ecol.*, 45: 94-98.
- Maregesi SM, Ngassapa OD, Pieters L, Vlietinck AJ (2007). Ethnopharmacological survey of the Bunda district, Tanzania: Plants used to treat infectious diseases. *J. Ethnopharmacol.*, 113(3): 457-470.
- Massei G, Hartley SE, Bacon PJ (2000). Chemical and Morphological Variation of Mediterranean Woody Evergreen Species: Do Plants Respond to Ungulate Browsing? *J. Veg. Sci.*, 11(1): 1-8.
- Mossholder K, Settoon R, Harris S, Armenakis A (1995). Measuring emotion in open-ended survey responses: An application of textual data analysis. *J. Manage.*, 21: 335-355.
- Mugisha-Kamatanesi M, Deng A, Ogendo J, Omolo E, Mihale M, Otim M, Buyungo J, Bett P (2008). Indigenous knowledge of field insect pests and their management around lake Victoria basin in Uganda. *Afr. J. Environ. Sci. Technol.*, 2: 342-348.
- Muhammad BY, Awaisu A (2008). The need for enhancement of research, development, and commercialization of natural medicinal products in Nigeria: Lessons from the Malaysian experience. *Afr. J. Trad., CAM*, 5(2): 120-130.
- Oerke E, Dehne HW (2004). Safeguarding production - losses in major crops and the role of crop protection. *Crop Protect.*, 23: 275-285.
- Oryemoriga H, Kakudidi EK Z, Katende AB, Bukenya ZR (1995). Preliminary ethnobotanical studies of the Rwenzori Mountain forest area in Bundibugyo district, Uganda. *Bothalia*, 25(1): 111-119.
- Palo RT (1984). Distribution of birch (*Betula* spp.), willow (*Salix* spp.), and poplar (*Populus* spp.) secondary metabolites and their potential role as chemical defense against herbivores. *J. Chem. Ecol.*, 10(3): 499-520.
- Pei SJ (2001). Ethnobotanical approaches of traditional medicine studies: Some experiences from Asia. *Pharm. Biol.*, 39: 74-79.
- Rates SM K (2001). Plants as source of drugs. *Toxicol.*, 39(5): 603-613.
- Schemske DW, Husband BC, Ruckelshaus MH, Goodwillie C, Parker IM, Bishop JG (1994). Evaluating Approaches to the Conservation of Rare and Endangered Plants. *Ecol.*, 75(3): 584-606.
- Schutte HR (1984). Secondary plant substances. Monoterpenes. *Prog. Bot.*, 46: 119-139.
- Scialabba Nadia (2000). Factors influencing organic agriculture policies with a focus on developing countries. IFOAM 2000 Scientific Conference, Basel, Switzerland, 28-31 August 2000. www.FAO.org/docs. Accessed on 23 June 2009.
- Sirikantaramas S, Yamazaki M, Saito K (2008). Mechanisms of resistance to self-produced toxic secondary metabolites in plants. *Phytochem. Rev.*, 7: 467-477.
- Ssegawa P, Kasenene JM (2007). Medicinal plant diversity and uses in the Sango bay area, Southern Uganda. *J. Ethnopharmacol.*, 113(3): 521-540.
- Swain T (1977). Secondary Compounds as Protective Agents. *Annu. Rev. Plant Physiol.*, 28(1): 479-501.
- Tabuti JRS, Dhillion SS, Lye K A (2003). Ethnoveterinary medicines for cattle (*Bos indicus*) in Bulamogi County, Uganda: plant species and mode of use. *J. Ethnopharmacol.*, 88(2-3): 279-286.
- Tenywa M, Isabirye M, Lal R, Lufafa A, Achan P (1999). Cultural practices and production constraints in smallholder banana-based cropping systems of Uganda's Lake Victoria Basin. *Afr. Crop Sci. J.*, 7: 613-623.
- Thacker JRM (2002). An introduction to arthropod pest control. Cambridge University Press, Cambridge. UK., p. 320.
- van Jaarsveld AS, Freitag S, Chown S L, Muller C, Koch S, Hull H, Bellamy C, Krüger M, Endrödy-Younga S, Mansell MW (1998). Biodiversity Assessment and Conservation strategies. *Science* 279(5359): 2106.
- Waako PJ, Katuura E, Smith P, Folb P (2007). East African medicinal plants as a source of lead compounds for the development of new antimalarial drugs. *Afr. J. Ecol.*, 45: 102-106.
- Walker K (2000). Cost-comparison of DDT and alternative insecticides for malaria control. *Med. Vet. Entomol.*, 14(4): 345-354.
- Walter KS, Gillett HJ (Eds) (1998). 1997 IUCN Red List of Threatened Plants. Gland, Switzerland and Cambridge, UK. The World Conservation Union, p. 862.

